

Exploring Benefits of a Be-Doped Carbon Nanocone for the Drug Delivery of 2-Thiouracil: Computational Study

Kun Harismah^{1,*}, Newsha Saeedi², Natthaporn Rattanapan³

¹ Department of Chemical Engineering, Faculty of Engineering, Universitas Muhammadiyah Surakarta, Surakarta, Indonesia; kun.harismah@ums.ac.id (K.H.);

² Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran; niyousha.saeedi@gmail.com (N.S.);

³ Faculty of Agricultural Technology, Phuket Rajabhat University, Muang, Phuket 83000, Thailand; rnatthaporn98@yahoo.com (N.R.);

* Correspondence: kun.harismah@ums.ac.id (K.H.);

Scopus Author ID 56982926300

Received: 19.09.2022; Accepted: 1.11.2022; Published: 3.01.2023

Abstract: By the importance of employing nanostructures in therapeutics, a model of beryllium (Be)-doped carbon nanocone (BNC) was explored in this work for the drug delivery of 2-thiouracil (2TU). Density functional theory (DFT) calculations were performed to recognize the stabilized structures in single and complex states. Six models of BNC-2TU complexes were obtained with different levels of interaction strengths, which were all reasonable levels of strength for the formation. In each model, a characteristic relaxation configuration of BNC and 2TU substances was observed with a deterministic role of defining strength levels and electronic features. Be...S, Be...O, and C...H were the types of observed interactions in the complexes with complementary roles of enhancing the strength of complexes. Localizations of molecular orbitals patterns were also dependent on the relaxation configurations of complexes; localization at both interacting sides or only one side was observed for the models. Indeed, obtaining such various features made the complexes flexible models for employing the desired functions and purposes. Formations of all models were suitable regarding the values of interaction energies; the obtained configurations and features could make them appropriate complexes for approaching desired functions such as drug delivery processes.

Keywords: nanocone; 2-thiouracil; adsorption; drug delivery; density functional theory.

© 2023 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

After the pioneering innovation of carbon nanotubes in 1991, several efforts have been dedicated to learning features of novel nanostructures to develop their desired applications [1-5]. A characteristic feature of high surface area made nanostructures suitable surfaces for working as an adsorbent in adsorption processes [6-10]. The pure carbon-based nanostructures were almost the first innovative nanostructures, but their modifications and heteroatomic compositions were achievable for these novel materials [11-15]. Existences of low-concentration atomic impurities could make atomic doped nanostructures with the suitability of heteroatomic regions for conducting the adsorption processes [16-20]. In this regard, several efforts have been dedicated to this time to make different atomic doped or molecular functionalized nanostructures regarding their suitability for desired purposes and processes [21-25]. Additionally, other geometrical shapes of nanostructures besides the original tubular

shape were found in the category of nanostructures, such as fullerene, graphene, and nanocone [26-30]. It should be noted that the shape of the nanostructure could make it a characteristic material for participating in the desired processes [31-35]. The synthetic processes have found the original nanocone as a conical shape with the carbon atoms compositions [36]. After that synthetic innovation, the nanocones have been topics of considerable research work up to now for developing their new applications [37-40]. The apex tip of nanocones is an important site for conducting communications with other external substances with specific features for developing applications, especially in interactions with other substances [41-45]. During the last three decades, several biomedical applications were developed for the nanostructures in single-standing nano-systems or combinations with other chemical and biochemical systems [46-50]. One of the major expected applications of nanostructures is their significant role in targeted drug delivery platforms [51-55]. In this regard, several efforts have been made to enhance the medical platforms with the assistance of nanostructures from the time of nanotechnology innovation up to now [56-60]. Accordingly, the benefits of a model of beryllium (Be) doped carbon nanocone (Figure 1) were explored in this work for employing in the drug delivery of 2-thiouracil (Figure 2).

2-Thiouracil (2TU) has been known as an anti-thyroid drug for several years by inhibiting the activity of the thyroid peroxidase enzyme [61-65]. Because of its important therapeutic features, several works were done to enhance this drug's efficiency for approaching better treatment levels [66-70]. Indeed, exploring novel treatments for known and unknown diseases is very important to maintain the balance of the human health system [71-75]. Accordingly, developments of both new drug substances and new drug treatments are essential for approaching therapeutic purposes [76-80]. As explored in the current work, combining a drug with nanostructures is one way of approaching an enhanced drug delivery platform [81-85]. To this aim, combinations of a Be-doped carbon nanocone and 2-thiouracil drug (Figure 3) were explored in this work to learn the benefits of formations of such complex systems for approaching a targeted drug delivery platform. To achieve the goals of this work, computational chemistry tools were employed to obtain structures and their related features [86-90].

2. Materials and Methods

This work was done using density functional theory (DFT) calculations at the B3LYP-D3/6-31G* computational level as implemented in the Gaussian program [91]. The calculations were done in two steps optimizing the geometries first and evaluating the features next. The models of single Be-doped carbon nanocone (BNC) (Figure 1) and 2-thiouracil (2TU) (Figure 2) were optimized to be prepared for participating in the interaction processes. Based on the examinations of all possible interaction situations of BNC and 2TU, six models of BNC-2TU complexes were obtained as designated by C1 to C6 (Figure 3). All the complex configurations were obtained by optimization calculations, and additional describing features were evaluated. As listed in Table 1, the strengths of interactions were analyzed using the evaluated features of the quantum theory of atoms in molecules (QTAIM) [92-94]. In this step, all involving interactions between the substances of complexes, as shown in Figure 3, were identified based on each energetic and distance magnitudes. Moreover, the differences in total energies of complexes and each of the interacting substances were obtained by the strengths of molecular interaction energies. Types of interactions and their distances (DIS), total electron density (RHO), Laplacian of electron density ($\text{DEL}^2\text{-RHO}$), energy density (H), and interaction

energy (E_I) were all summarized in Table 1. It is worth to note the calculated values of basis set superposition error (BSSE) were almost negligible to be considered in the evaluations of values of E_I . Next, the contents of Table 2 were provided by the values of molecular orbitals energies emphasizing the values of the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO) and their related features, such as energy gap (E_G) and chemical hardness (CH). Besides the evaluated quantities, graphical representations of HOMO and LUMO were exhibited in each of Figures 1-3 for the optimized geometries of the obtained models to show localization concentrations of molecular orbitals on the molecular sites. As a consequence, the models were available by the optimized geometries and the evaluated describing features for exploring the benefits of BNC for the drug delivery of 2TU.




Figure 1. The optimized model and HOMO-LUMO representations of single BNC.




Figure 2. The optimized model and HOMO-LUMO representations of single 2TU.

3. Results and Discussion

Exploring the benefits of employing a beryllium (Be) doped carbon nanocone (BNC) for the drug delivery of 2-thiouracil (2TU) was investigated in this work by means of performing DFT calculations. As shown in Figures 1 and 2, the single models were optimized to be prepared for participating in interactions with each other. For the BNC model, one Be atom was substituted at the tip of the nanocone to provide an interacting site for the structure. On the other hand, all possibilities of interactions of 2TU through its atomic sites were examined for participating in interactions with the adsorbent. Accordingly, six models of BNC-2TU complexes were obtained by performing optimization calculations as shown in Figure 3; C1-C6 complexes. It is worth mentioning that the obtained BNC-2TU complexes were obtained by examining all possibilities of interactions between BNC and 2TU substances, in which six complexes were obtained. To make a deep analysis of the complexes, details of interactions were investigated by performing additional QTAIM analysis on the optimized BNC-2TU complexes.




Figure 3. The optimized models and HOMO-LUMO (H and L) representations of complexes of BNC-2TU.

Table 1. Interactions features.¹

BNC-2TU	INT	DIS (Å)	RHO (au)	DEL ² -RHO (au)	H (au)	E ₁ (kcal/mol)
C1	Be...S	2.014	0.074	0.188	-0.051	-50.129
	C...H	1.451	0.068	0.037	-0.022	
C2	Be...S	2.006	0.069	0.204	-0.019	-42.344
	C...H	1.748	0.058	0.069	-0.011	
C3	Be...O	1.546	0.094	0.719	-0.064	-50.968
	C...H	1.745	0.061	0.069	-0.012	
C4	Be...O	1.579	0.082	0.633	-0.019	-45.581
	C...H	2.599	0.013	0.033	-0.011	
C5	Be...C	1.885	0.056	0.179	-0.014	-17.626
	H...C	2.595	0.009	0.025	-0.001	
C6	H...C	2.595	0.009	0.025	-0.001	-2.672

¹The models were shown in Figure 3.

The involved interactions of Figure 3 were known by the obtained QTAIM results, in which their detailed quantities were summarized in Table 1. A quick achievement of the obtained results is that the Be-doped region worked as an interaction environment manager to manage interactions between BNC and 2TU substances. In all complex cases, the models communicated with each other through the Be-doped atom directly or its neighboring carbon atom indirectly. Additionally, the idea of Be-doping of nanocone was affirmed by a significant contribution of Be atom of BNC substance to interactions with the 2TU substance. In this regard, the models were stabilized, and their features were investigated in terms of geometries and energies to show the benefits of formations of such BNC-2TU complexes. To this point, the models of complexes were also comparable to each other regarding their priority of formation. As described in Table 1, the complex models of Figure 3 were in different levels of strength of interactions, and different configurations were obtained. The important point is that the interactions were all in the physical mode of non-covalent type, and their strengths were reasonable. Another important point is that the existence of C...H interaction was observed in the complexes. Additionally, different configurations and their corresponding strengths showed the importance of examining all interacting configurations to approach complex formations. Based on the obtained values of EI, the strength order of interacting complexes was found to be C3 > C1 > C4 > C2 > C5 > C6. In this regard, the obtained strengths showed the importance of the relaxation configuration of two counterparts towards each other and the characteristic role of their involving interactions. Accordingly, both of Be...S and Be...O interactions were stronger by the existence of additional C...H interactions. As a consequence, formations of BNC-2TU complexes were found suitable for approaching the role of adsorbent for the BNC substance towards the 2TU substance.

Table 2. Molecular orbitals features.¹

Model	HOMO (eV)	LUMO (eV)	E _G (eV)	C _H (eV)
BNC	-5.241	-2.709	2.532	1.266
2TU	-6.315	-1.604	4.711	2.356
C1	-5.060	-2.246	2.814	1.407
C2	-4.536	-2.413	2.123	1.061
C3	-4.491	-2.920	1.571	0.785
C4	-4.067	-3.424	0.644	0.322
C5	-4.757	-3.254	1.502	0.751
C6	-5.476	-3.159	2.316	1.158

¹The models are shown in Figures 1-3.

Molecular orbitals of the optimized modes were exhibited in Figures 1-3, and their quantities were summarized in Table 2 based on the consideration of HOMO and LUMO levels. Indeed, the frontier molecular orbitals play important roles in determining the electronic features of molecular models, especially within their participation in electronic transferring processes. To this aim, such features were evaluated for the models of this work to describe their electronic-related properties. As shown in Figure 1, a significant concentration of LUMO at the Be-doped region of BNC made it a suitable region for participating in interactions through the lone pairs of electrons of S and O atoms of 2TU. Accordingly, the models of C1 to C6 showed different configurations of HOMO and LUMO patterns based on different configurations of relaxations of two interacting BNC and 2TU substances. In the cases of C2, C3, C4, and C5, patterns of HOMO and LUMO were separated with the HOMO localization at the BNC part and the LUMO localization at the 2TU part. In the case of C1, both of HOMO and LUMO were localized at the BNC part. And in the case of C6, a model of pattern localization similar to the original single BNC was observed. In this regard, the models were

distinguished based on their electronic features with deterministic roles for their future applications in the drug delivery processes. In addition to these graphical representations, the evaluated quantities of Table 2 showed different features between single and complex models. The levels of HOMO and LUMO were significantly changed from single to complex states, besides observing significant variations among the complex models. Accordingly, the values of E_G and C_H were changed regarding the changes in HOMO and LUMO levels. It is obvious that the formation of BNC-2TU complexes had impacts on the electronic features, in which the model of configuration relaxation of BNC and 2TU towards each other was dominant for observing the molecular orbital-based features. In this regard, the desired function could determine the advantage of each complex formation for approaching the goal.

4. Conclusions

To summarize this work's achievements, an important point is the reasonable formation of BNC-2TU complexes regarding the obtained levels of interaction strengths. Additionally, six possible configurations were obtained for the complexes regarding the relaxation of each of the BNC and 2TU substances towards each other. The Be-doped region was a dominant site of interactions for the BNC substance to manage formations of complexes. Regarding the graphical and quantitative features of molecular orbitals, the impacts of such relaxation configurations of complexes were observed for the models. Localizations of each of the HOMO and LUMO patterns at each molecular side or both molecular sides were determined by changes in relaxation configurations. Accordingly, the models were found flexible for approaching the desired function, such as conducting the drug delivery processes.

Funding

This research received no external funding.

Acknowledgments

This research has no acknowledgment.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Li, Y. Carbon nanotube research in its 30th year. *ACS Nano* **2021**, *15*, 9197–9200, <https://doi.org/10.1021/acsnano.1c04972>.
2. Bale, V.; Katreddi, H. Synthesis, characterization and catalytic activity of zinc oxide nanoparticles functionalized with metallo-thiosemicarbazones. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 159–173, <https://doi.org/10.26655/AJNANOMAT.2022.2.7>.
3. Ayoub, H.; Khairy, M.; Rashwan, F.; Abdel-Hafez, H. Nanomaterial-based agrochemicals new avenue for sustainable agriculture: a short review. *Journal of Chemical Reviews* **2022**, *4*, 191–199, <https://doi.org/10.22034/jcr.2022.336130.1163>.
4. Ebadian, B.; Fathi, A.; Beiranvand, N. Investigation of the effect of bonding factors on strength of porcelain bond to soft metal alloys after application of thermal cycle. *Dental Research Journal* **2022**, *19*, 91, <https://doi.org/10.4103/1735-3327.359328>.
5. Hajinasiri, R.; Esmaeili Jadidi, M. Synthesis of ZnO nanoparticles via flaxseed aqueous extract. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 101–108, <https://doi.org/10.22034/jaoc.2022.154805>.

6. Hassani, M.; Zeeb, M.; Monzavi, A.; Khodadadi, Z.; Kalaei, M. Response surface modeling and optimization of microbial fuel cells with surface-modified graphite anode electrode by ZSM-5 nanocatalyst functionalized. *Chemical Methodologies* **2022**, *6*, 253–268, <https://doi.org/10.22034/chemm.2022.324312.1425>.
7. Odogu Ankoro, N.; Ngouateu Lekene, R.; Ndi, J.; Kouotou, D.; Ngomo, H.; Ketcha, J. Highly microporous activated carbons from *Mangifera indica* residues: optimization of preparation conditions using response surface methodology. *Asian Journal of Green Chemistry* **2022**, *6*, 1–13, <https://doi.org/10.22034/ajgc.2022.1.1>.
8. Janighorban, M.; Rasouli, N.; Sohrabi, N.; Ghaedi, M. Use of central composite design and surface modeling for cadmium (ii) ions removal from aqueous environment using novel modified layer double hydroxide. *Asian Journal of Green Chemistry* **2021**, *5*, 151–175, <https://doi.org/10.22034/ajgc.2021.118922>.
9. Udeh, J.; Nkele, A.; Obodo, R.; Nwodo, I.; Chime, C.; Nwanya, A.; Ezema, F.; Maaza, M. Investigating the properties of cobalt phosphate nanoparticles synthesized by co-precipitation method. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 22–35, <https://doi.org/10.26655/AJNANOMAT.2022.1.3>.
10. Fathpour, K.; Ahmadabadi, M.N.; Atash, R.; Fathi, A. Effect of different surface treatment methods on the shear bond strength of resin composite/zirconia for intra-oral repair of zirconia restorations. *European Journal of Dentistry* **2022**, *in press*, <http://doi.org/10.1055/s-0042-1756475>.
11. Alizadeh, K.; Khaledyan, E.; Mansourpanah, Y. Novel modified magnetic mesoporous silica for rapid and efficient removal of methylene blue dye from aqueous media. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 224–234, <https://doi.org/10.22034/jaoc.2022.155004>.
12. Dessie, Y.; Tadesse, S. A review on advancements of nanocomposites as efficient anode modifier catalyst for microbial fuel cell performance improvement. *Journal of Chemical Reviews* **2021**, *3*, 320–344, <https://doi.org/10.22034/jcr.2021.314327.1128>.
13. Shete, R.; Fernandes, P.; Borhade, B.; Pawar, A.; Sonawane, M.; warude, N. Review of cobalt oxide nanoparticles: green synthesis, biomedical applications, and toxicity studies. *Journal of Chemical Reviews* **2022**, *4*, 331–345, <https://doi.org/10.22034/jcr.2022.342398.1172>.
14. Baghernejad, B. Preparation of polyhydroquinolines in the presence of nano cerium (IV)oxide/zinc oxide as an efficient catalyst. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 74–80, <https://doi.org/10.22034/jaoc.2022.340645.1054>.
15. Patel, K.; Chandankar, S.; Mahajan, H. Synthesis and characterization of curcumin nanoparticles by hydrothermal method. *Asian Journal of Nanosciences and Materials* **2021**, *4*, 321–330, http://www.ajnanomat.com/article_139267.html.
16. Tosan, F.; Rahnama, N.; Sakhaei, D.; Fathi, A.; Yari, A. Effects of doping metal nanoparticles in hydroxyapatite in improving the physical and chemical properties of dental implants. *Nanomedicine Research Journal* **2021**, *6*, 327–336, <https://doi.org/10.22034/nmrj.2021.04.002>.
17. Sengar, M.; Saxena, S.; Lakhani, A.; Satsangee, S. Allium based green route synthesized silver nanoparticles for removal of polycyclic aromatic hydrocarbons. *Asian Journal of Green Chemistry* **2021**, *5*, 206–218, <https://doi.org/10.22034/ajgc.2021.271486.1293>.
18. Shayegan, H.; Safari Fard, V.; Taherkhani, H.; Rezvani, M. Efficient removal of cobalt(II) ion from aqueous solution using amide-functionalized metal-organic framework. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 109–118, <https://doi.org/10.22034/jaoc.2022.154718>.
19. Sheikholeslami-Farahani, F. Amine functionalized SiO₂@Fe₃O₄ as a green and reusable magnetic nanoparticles system for the synthesis of Knoevenagel condensation in water. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 132–143, <https://doi.org/10.26655/AJNANOMAT.2022.2.5>.
20. Baghernejad, B.; Hojjati Taromsari, S. Aqueous media preparation of 2-amino-4H-benzopyran derivatives using cerium oxide nanoparticles as a recyclable catalyst. *Asian Journal of Green Chemistry* **2022**, *6*, 194–202, <https://doi.org/10.22034/ajgc.2022.3.2>.
21. Golipour-Chobar, E.; Salimi, F.; Ebrahimzadeh-Rajaei, G. Sensing of lomustine drug by pure and doped C48 nanoclusters: DFT calculations. *Chemical Methodologies* **2022**, *6*, 790–800, <https://doi.org/10.22034/chemm.2022.344895.1555>.
22. Baghernejad, B.; Khoshnud Gilakejan, S. A new strategy for the synthesis of dihydropyrano[c]chromene using zinc oxide/copper oxide as a nano and efficient catalyst. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 1–10, <https://doi.org/10.26655/AJNANOMAT.2022.1.1>.
23. Safari Fard, V.; davoudabadi farahani, Y. An amine/imine functionalized microporous MOF as a new fluorescent probe exhibiting selective sensing of Fe³⁺ and Al³⁺ over mixed metal ions. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 180–194, <https://doi.org/10.22034/jaoc.2022.154981>.

24. Kavade, R.; Khanapure, R.; Gawali, U.; Patil, A.; Patil, S. Degradation of Methyl orange under visible light by ZnO-polyaniline nanocomposites. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 89–100, <https://doi.org/10.22034/jaoc.2022.349558.1056>.
25. Hasanpour, F.; Taei, M.; Fouladgar, M.; Salehi, M. Au nano dendrites/composition optimized Nd-doped cobalt oxide as an efficient electrocatalyst for ethanol oxidation. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 203–211, <https://doi.org/10.22034/jaoc.2022.154984>.
26. Jasim, S.A.; Kzar, H.H.; Jalil, A.T.; Kadhim, M.M.; Mahmoud, M.Z.; Al-Gazally, M.E.; Nasser, H.A.; Ahmadi, Z. DFT investigation of BN, AlN, and SiC fullerene sensors for arsine gas detection and removal. *Main Group Chemistry* **2022**, *21*, 513–521, <https://doi.org/10.3233/MGC-210145>.
27. Hamidi Zare, S.; Ghorayshi Nejad, M.; Saberi, A.; Rostami, E. A highly efficient and sustainable synthesis of 1-[(1,3-thiazol-2-ylamino) methyl]-2-naphthols under solvent-free conditions using graphene oxide substituted ethane sulfonic acid catalyst. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 211–224, http://www.ajnanomat.com/article_155115.html.
28. Hakimi Saryazdi, F.; Golrasan, E.; Heidari, S. Nano particles graphene oxid: a green and effective catalyst for synthesis of pyrazoles. *Asian Journal of Green Chemistry* **2021**, *5*, 325–334, <https://doi.org/10.22034/ajgc.2021.273614.1295>.
29. Sun, Z.; Wang, S.; Xiong, H.; Wu, K.; Shi, J. Optimal nanocone geometry for water flow. *AIChE Journal* **2022**, *68*, e17543, <https://doi.org/10.1002/aic.17543>.
30. Shoyama, K.; Würthner, F. Synthesis of a carbon nanocone by cascade annulation. *Journal of the American Chemical Society* **2019**, *141*, 13008–13012, <https://doi.org/10.1021/jacs.9b06617>.
31. Tabar Maleki, S.; Sadati, S. Investigation of magnetic properties of Fe₃O₄/Halloysite nanotube/polypyrrole core-shell nanocomposite and its stability in the acidic environment. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 98–108, <https://doi.org/10.26655/AJNANOMAT.2022.2.2>.
32. Sukmawati, A.; Da'i, M.; Yuliani, R.; Anggraeni, S.N.; Wahyuningsih, D. Characterization of chitosan nanoparticle containing combination doxorubicin and curcumin analogue. *Advanced Science Letters* **2017**, *23*, 12486–12488, <https://doi.org/10.1166/asl.2017.10798>.
33. Baghernejad, B.; Samaie, R. Nano CeO₂ as a new green and recyclable catalyst for the synthesis of 2-aryl Benzoxazole. *Asian Journal of Nanosciences and Materials* **2021**, *4*, 255–262, http://www.ajnanomat.com/article_132310.html.
34. Harismah, K.; Mirzaei, M.; Da'i, M.; Roshandel, Z.; Salarrezaei, E. In silico investigation of nanocarbon biosensors for diagnosis of COVID-19. *Eurasian Chemical Communications* **2021**, *3*, 95–102, <https://doi.org/10.22034/ecc.2021.267226.1120>.
35. Sukmawati, A.; Utami, W.; Yuliani, R.; Da'i, M.; Nafarin, A. Effect of tween 80 on nanoparticle preparation of modified chitosan for targeted delivery of combination doxorubicin and curcumin analogue. *IOP Conference Series: Materials Science and Engineering* **2018**, *311*, 012024, <https://doi.org/10.1088/1757-899X/311/1/012024>.
36. Yang, N.; Zhang, G.; Li, B. Carbon nanocone: a promising thermal rectifier. *Applied Physics Letters* **2008**, *93*, 243111, <https://doi.org/10.1063/1.3049603>.
37. Fu, H.; Bao, H.; Zhang, H.; Zhao, Q.; Zhou, L.; Zhu, S.; Wei, Y.; Li, Y.; Cai, W. Quantitative surface-enhanced raman spectroscopy for field detections based on structurally homogeneous silver-coated silicon nanocone arrays. *ACS Omega* **2021**, *6*, 18928–18938, <https://doi.org/10.1021/acsomega.1c02179>.
38. EL-Barbary AA, Alkhateeb MA. DFT study of Se-doped nanocones as highly efficient hydrogen storage carrier. *Graphene* **2021**, *10*, 49–60, <https://doi.org/10.4236/graphene.2021.104004>.
39. García-Hernández, E.; Flores-López, A.; García-Contreras, M.A.; Palomino-Asencio, L.; Catarino-Centeno, R. Theoretical analysis of the uptake of CO, CO₂, and NO₂ on pristine and BN-doped carbon nanocones. *Chemical Physics Letters* **2022**, *795*, 139531, <https://doi.org/10.1016/j.cplett.2022.139531>.
40. Koam, A.N.; Ahmad, A.; Azeem, M.; Nadeem, M.F.; Bounds on the partition dimension of one pentagonal carbon nanocone structure. *Arabian Journal of Chemistry* **2022**, *15*, 103923, <https://doi.org/10.1016/j.arabjc.2022.103923>.
41. Kamel Attar Kar, M.H.; Yousefi, M. Interaction of a conical carbon scaffold with the thio-substituted model of fluorouracil towards approaching the drug delivery purposes. *Main Group Chemistry* **2022**, *21*, 725–735, <https://doi.org/10.3233/MGC-210174>.
42. Al-Haideri, L.M.; Cakmak, N.; Mahdi Aal-Shabeeb, K.H.; Dhumad, A.M. DFT calculations of plutonium-doped conical nanocarbons: Exploring structural and electronic features. *Main Group Chemistry* **2022**, *in press*, <https://doi.org/10.3233/MGC-220023>.

43. Yodsin, N.; Sakagami, H.; Udagawa, T.; Ishimoto, T.; Jungsuttiwong, S.; Tachikawa, M. Metal-doped carbon nanocones as highly efficient catalysts for hydrogen storage: nuclear quantum effect on hydrogen spillover mechanism. *Molecular Catalysis* **2021**, *504*, 111486, <https://doi.org/10.1016/j.mcat.2021.111486>.
44. Kose, A.; Yuksel, N.; Fellah, M.F. Hydrogen adsorption on Ni doped carbon nanocone. *Diamond and Related Materials* **2022**, *124*, 108921, <https://doi.org/10.1016/j.diamond.2022.108921>.
45. Liu, P.; Chen, B.; Liang, C.; Yao, W.; Cui, Y.; Hu, S.; Zou, P.; Zhang, H.; Fan, H.J.; Yang, C. Tip-enhanced electric field: a new mechanism promoting mass transfer in oxygen evolution reactions. *Advanced Materials* **2021**, *33*, 2007377, <https://doi.org/10.1002/adma.202007377>.
46. Saliminasab, M.; Jabbari, H.; Farahmand, H.; Asadi, M.; Soleimani, M.; Fathi, A. Study of antibacterial performance of synthesized silver nanoparticles on *Streptococcus mutans* bacteria. *Nanomedicine Research Journal* **2022**, *7*, 391–396, <https://doi.org/10.22034/nmrj.2022.04.010>.
47. Baghernejad, B.; Alikhani, M. Nano-cerium oxide/aluminum oxide as an efficient catalyst for the synthesis of xanthene derivatives as potential antiviral and anti-inflammatory agents. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 155–162, <https://doi.org/10.22034/jaoc.2022.154819>.
48. Mortezagholi, B.; Movahed, E.; Fathi, A.; Soleimani, M.; Forutan Mirhosseini, A.; Zeini, N.; Khatami, M.; Naderifar, M.; Abedi Kiasari, B.; Zareanshahraki, M. Plant-mediated synthesis of silver-doped zinc oxide nanoparticles and evaluation of their antimicrobial activity against bacteria cause tooth decay. *Microscopy Research and Technique* **2022**, *85*, 3553, <https://doi.org/10.1002/jemt.24207>.
49. Alijani, H.Q.; Fathi, A.; Amin, H.I.; Lima Nobre, M.A.; Akbarizadeh, M.R.; Khatami, M.; Jalil, A.T.; Naderifar, M.; Dehkordi, F.S.; Shafiee, A. Biosynthesis of core-shell α -Fe₂O₃@ Au nanotruffles and their biomedical applications. *Biomass Conversion and Biorefinery* **2022**, *in press*, <https://doi.org/10.1007/s13399-022-03561-3>.
50. Alinezhad, H.; Hajiabbas Tabar Amiri, P.; Mohseni Tavakkoli, S.; Muhiebes, R.; Fakri Mustafa, Y. Progressive types of Fe₃O₄ nanoparticles and their hybrids as catalysts. *Journal of Chemical Reviews* **2022**, *4*, 288–312, <https://doi.org/10.22034/jcr.2022.325255.1137>.
51. Abdul Ahad, H.; Chinthaginjala, H.; Yaparla, S.; Snehitha, B.; Tanuja, M.; Srinivasa Sainath, K. A desk-top literature for research on gas engendering and low-density floating drug delivery systems. *Journal of Chemical Reviews* **2022**, *4*, 147–155, <https://doi.org/10.22034/jcr.2022.332618.1153>.
52. Maghsoudi, S.; Hosseini, S.; Ravandi, S. A review on phospholipid and liposome carriers: synthetic methods and their applications in drug delivery. *Journal of Chemical Reviews* **2022**, *4*, 346–363, <https://doi.org/10.22034/jcr.2022.355104.1182>.
53. Yaraghi, A.; Ozkendir, O.M.; Mirzaei, M. DFT studies of 5-fluorouracil tautomers on a silicon graphene nanosheet. *Superlattices and Microstructures* **2015**, *85*, 784–788, <https://doi.org/10.1016/j.spmi.2015.05.053>.
54. Darougari, H.; Rezaei-Sameti, M. The drug delivery appraisal of Cu and Ni decorated B12N12 nanocage for an 8-hydroxyquinoline drug: A DFT and TD-DFT computational study. *Asian Journal of Nanosciences and Materials* **2022**, *5*, 196–210, http://www.ajnanomat.com/article_154625.html.
55. Hosseini, S.; Shojaie, F.; Afzali, D. Analysis of structural, and electronic properties of clopidogrel drug adsorption on armchair (5, 5) single-walled carbon nanotube. *Asian Journal of Nanosciences and Materials* **2021**, *4*, 15–30, <https://doi.org/10.26655/AJNANOMAT.2021.1.2>.
56. Tavakoli, F.; Shafiei, H.; Ghasemikhah, R. Kinetic and thermodynamics analysis: effect of eudragit polymer as drug release controller in electrospun nanofibers. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 235–243, <https://doi.org/10.22034/jaoc.2022.155078>.
57. Zilabi, S.; Shareei, M.; Bozorgian, A.; Ahmadpour, A.; Esmaeil, E. A review on nanoparticle application as an additive in lubricants. *Advanced Journal of Chemistry-Section B* **2022**, *4*, 209–221, <https://doi.org/10.22034/ajcb.2022.353097.1125>.
58. Aminian, A.; Fathi, A.; Gerami, M.H.; Arsan, M.; Forutan Mirhosseini, A.; Torabizadeh, S.A. Nanoparticles to overcome bacterial resistance in orthopedic and dental implants. *Nanomedicine Research Journal* **2022**, *7*, 107–123, <https://doi.org/10.22034/nmrj.2022.02.001>.
59. Ahmad, F.; Mehmood, M. A critical review of photocatalytic degradation of organophosphorus pesticide “Parathion” by different mixed metal oxides. *Advanced Journal of Chemistry-Section A* **2022**, *5*, 287–310, <https://doi.org/10.22034/ajca.2022.357664.1324>.
60. Alizadeh, S.; Nazari, Z. Amphetamine, methamphetamine, morphine @ AuNPs kit based on PARAFAC. *Advanced Journal of Chemistry-Section A* **2022**, *5*, 253–262, <https://doi.org/10.22034/ajca.2022.345350.1318>.

61. Lara, N.L.; Silva Jr, V.A.; Chiarini-Garcia, H.; Garcia, S.K.; Debeljuk, L.; Hess, R.A.; França, L.R. Hypothyroidism induced by postnatal PTU (6-n-propyl-2-thiouracil) treatment decreases Sertoli cell number and spermatogenic efficiency in sexually mature pigs. *General and Comparative Endocrinology* **2020**, *299*, 113593, <https://doi.org/10.1016/j.ygcen.2020.113593>.
62. Lee, J.H.; Koo, J.S.; Roh, S.S.; Park, J.H.; Seo, B.I. The effects of astragali radix on hypothyroidism rat model induced by 6-propyl-2-thiouracil (PTU). *The Korea Journal of Herbology* **2019**, *34*, 45–53, <https://doi.org/10.6116/kjh.2019.34.3.45>.
63. Awad, S.M.; Zohny, Y.M.; Ali, S.A.; Mahgoub, S.; Said, A.M. Design, synthesis, molecular modeling, and biological evaluation of novel thiouracil derivatives as potential anti-thyroid agents. *Molecules* **2018**, *23*, 2913, <https://doi.org/10.3390/molecules23112913>.
64. Khalil, N.Y.; AlRabiah, H.K.; Almousa, M.S.; Bari, A.; Alkahtani, H.M. Thiouracil: comprehensive profile. *Profiles of Drug Substances, Excipients and Related Methodology* **2019**, *44*, 293–331, <https://doi.org/10.1016/bs.podrm.2018.11.006>.
65. Léger, J.; Delcour, C.; Carel, J.C. Fetal and neonatal thyroid dysfunction. *The Journal of Clinical Endocrinology & Metabolism* **2022**, *107*, 836–846, <https://doi.org/10.1210/clinem/dgab747>.
66. Alcolea Palafox, M.; Franklin Benial, A.M.K.; Rastogi, V. Biomolecules of 2-thiouracil, 4-thiouracil and 2,4-dithiouracil: a DFT study of the hydration, molecular docking and effect in DNA:RNA microhelices. *International Journal of Molecular Sciences* **2019**, *20*, 3477, <https://doi.org/10.3390/ijms20143477>.
67. Hussein, T.; Ahamad, M. Synthesis of C- glycoside analogs containing thiouracil derivatives and evaluated as antibacterial, antifungal and antioxidant activities. *Eurasian Chemical Communications* **2021**, *3*, 584–589, <https://doi.org/10.22034/ecc.2021.293403.1197>.
68. Woźniak, B.; Witek, S.; Matraszek-Żuchowska, I.; Sell, B.; Posyniak, A. The effect of diet enriched with rapeseed meal on endogenous thiouracil contents in urine of calves. *Research in Veterinary Science* **2021**, *136*, 192–197, <https://doi.org/10.1016/j.rvsc.2020.10.021>.
69. Skrobanska, M.; Zabizsak, M.; Taras-Goslinska, K.; Nowak, M.; Kaczmarek, M.T.; Frymark, J.; Michalska, D.; Jastrzab, R. Potentiometric and spectroscopic studies of the complex formation in the 6-methyl-2-thiouracil and copper (ii) ion system. *Polyhedron* **2022**, *223*, 115964, <https://doi.org/10.1016/j.poly.2022.115964>.
70. Cao, Y.; Rajhi, A.A.; Yousefi, M.; Ahmadi, R. Investigating thiouracil adsorption by an iron-doped carbon particle: analyzing structural, electronic, and QTAIM features. *Journal of Molecular Structure* **2022**, *1250*, 131885, <https://doi.org/10.1016/j.molstruc.2021.131885>.
71. Hatami, A. Preparation, description and evaluation of the lethality acid loaded liposomal nanoparticles against in vitro colon and liver cancer. *Journal of Chemical Reviews* **2021**, *3*, 121–133, <https://doi.org/10.22034/jcr.2021.283609.1109>.
72. Sadeghi, P.M.; Rabiee, M.; Darestani, N.G.; Alesahab, F.; Zeinalkhani, F. Short term results of stapled versus conventional hemorrhoidectomy within 1 year follow-up. *International Journal of Burns and Trauma* **2021**, *11*, 69–74, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8012873>.
73. Kiasari, B.A.; Abbasi, A.; Darestani, N.G.; Adabi, N.; Moradian, A.; Yazdani, Y.; Hosseini, G.S.; Gholami, N.; Janati, S. Combination therapy with nivolumab (anti-PD-1 monoclonal antibody): a new era in tumor immunotherapy. *International Immunopharmacology* **2022**, *113*, 109365, <https://doi.org/10.1016/j.intimp.2022.109365>.
74. Fathi, A.; Rismanchian, M.; Dezaki, S.N. Effectiveness of different antimicrobial agents on malodor prevention in two-stage dental implants: a double-blinded randomized clinical trial. *European Journal of Dentistry* **2022**, *in press*, <https://doi.org/10.1055/s-0042-1747954>.
75. Khamisi, N.; Fathi, A.; Yari, A. Antimicrobial resistance of staphylococcus aureus isolated from dental plaques. *Academic Journal of Health Sciences: Medicina Balear* **2022**, *37*, 136–140, <https://doi.org/10.3306/AJHS.2022.37.01.136>.
76. Karbasizade, S.; Ghorbani, F.; Darestani, N.G.; Mansouri-Tehrani, M.M.; Kazemi, A.H. Comparison of therapeutic effects of statins and Aloe vera mouthwash on chemotherapy induced oral mucositis. *International Journal of Physiology, Pathophysiology and Pharmacology* **2021**, *13*, 110–116, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8446774>.
77. Hudiyawati, D.; Syafitry, W. Effectiveness of physical and psychological treatment for cancer-related fatigue: systematic review. *Jurnal Kesehatan* **2021**, *14*, 195–211, <https://doi.org/10.23917/jk.v14i2.15596>.
78. Babak, A.; Motamedi, N.; Mousavi, S.Z.; Darestani, N.G. Effects of mindfulness-based stress reduction on blood pressure, mental health, and quality of life in hypertensive adult women: a randomized clinical trial

- study. *The Journal of Tehran University Heart Center* **2022**, *17*, 127–133, <https://doi.org/10.18502/jthc.v17i3.10845>.
79. Azizian, M.; Darestani, N.G.; Aliabadi, A.; Afzali, M.; Tavoosi, N.; Fosouli, M.; Khataei, J.; Aali, H.; Nourian, S.M. Predictive value of number and volume of demyelinating plaques in treatment response in patients with multiple sclerosis treated with INF-B. *American Journal of Neurodegenerative Disease* **2022**, *11*, 10–16, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9123433>.
80. Saroyo, H.; Saputri, N.F. Cytotoxicity of mangrove leaves (rhizophora) ethanolic extract on cancer cells. *Journal of Nutraceuticals and Herbal Medicine* **2021**, *4*, 43–52, <https://doi.org/10.23917/jnhm.v4i1.15657>.
81. Qiao, L.; Yang, H.; Gao, S.; Li, L.; Fu, X.; Wei, Q. Research progress on self-assembled nanodrug delivery systems. *Journal of Materials Chemistry B* **2022**, *10*, 1908–1922, <https://doi.org/10.1039/D1TB02470A>.
82. Milani Fard, A.; Milani Fard, M. Modeling drug release. *Eurasian Journal of Science and Technology* **2022**, *2*, 14–31, <https://doi.org/10.22034/EJST.2022.1.2>.
83. Al Samarrai, E.; Alwan, L.; Al-Haddad, S.; Al Samarrai, M.; Al-Obaidi, M.; Al Samarrai, O. Spectrophotometric determination of phenobarbital in pharmaceutical preparation using gold nanoparticles. *Eurasian Chemical Communications* **2022**, *4*, 812–825, <https://doi.org/10.22034/ecc.2022.329806.1329>.
84. Milani Fard, M.; Milani Fard, A. Investigation of drug release from a biodegradable biphasic polymer system. *Eurasian Journal of Science and Technology* **2022**, *2*, 1–13, <https://doi.org/10.22034/EJST.2022.1.1>.
85. Hosseini, S.; Naimi-Jamal, M.; Hassani, M. Preparation and characterization of mebeverine hydrochloride niosomes as controlled release drug delivery system. *Chemical Methodologies* **2022**, *6*, 591–603, <https://doi.org/10.22034/chemm.2022.337717.1482>.
86. Venkatesh, G.; Sheena mary, Y.; Shymamary, Y.; Palanisamy, V.; Govindaraju, M. Quantum chemical and molecular docking studies of some phenothiazine derivatives. *Journal of Applied Organometallic Chemistry* **2021**, *1*, 148–158, <https://doi.org/10.22034/jaoc.2021.303059.1033>.
87. Merajoddin, M.; Piri, S.; Mokarian, Z.; Piri, F. Density functional theory study of benzoic acid decarboxylation. *Asian Journal of Green Chemistry* **2021**, *5*, 335–342, <https://doi.org/10.22034/ajgc.2021.302161.1310>.
88. Mohamed, H.; Hamza, Z.; Nagdy, A.; El-Mageed, H. Computational studies and DFT calculations of synthesized triazolo pyrimidine derivatives: a review. *Journal of Chemical Reviews* **2022**, *4*, 156–190, <https://doi.org/10.22034/jcr.2022.325439.1138>.
89. Farhami, N. A computational study of thiophene adsorption on boron nitride nanotube. *Journal of Applied Organometallic Chemistry* **2022**, *2*, 163–172, <https://doi.org/10.22034/jaoc.2022.154821>.
90. Adole, V. Computational chemistry approach for the investigation of structural, electronic, chemical and quantum chemical facets of twelve biginelli adducts. *Journal of Applied Organometallic Chemistry* **2021**, *1*, 29–40, <https://doi.org/10.22034/jaoc.2021.278598.1009>.
91. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R. Gaussian 09 program. *Gaussian Inc. Wallingford, CT*, **2009**. https://s3-eu-west-1.amazonaws.com/pstorage-acsc-6854636/3955375/jp500312r_si_001.pdf.
92. Pour Karim, S.; Ahmadi, R.; Yousefi, M.; Kalateh, K.; Zarei, G. Interaction of graphene with amoxicillin antibiotic by in silico study. *Chemical Methodologies* **2022**, *6*, 861–871, <https://doi.org/10.22034/chemm.2022.347571.1560>.
93. Srivastava, A.K. Ionization of NO by superhalogens: DFT and QTAIM approaches. *Main Group Chemistry* **2021**, *20*, 33–40, <https://doi.org/10.3233/mgc-210004>.
94. Bader, R.F.; Matta, C.F. Atomic charges are measurable quantum expectation values: a rebuttal of criticisms of QTAIM charges. *The Journal of Physical Chemistry A* **2004**, *108*, 8385–8394, <https://doi.org/10.1021/jp0482666>.